

C1 of binding to the α_4 subunit of VLA-4, in an amount effective to treat diabetes.

C2 ~~8-12~~. (Amended) A method according to claim ~~11~~ ¹/~~10~~, wherein the soluble VCAM-1 polypeptide[s] comprise a VCAM-IgG fusion.

~~3~~ ¹/~~13~~. (Amended) A method according to claim ~~11~~ ¹/~~10~~, wherein the composition is administered in an amount effective to provide a plasma level of a soluble VCAM-1 polypeptide[s] in the mammal of at least 10-20 $\mu\text{g/ml}$ over a period of 1-14 days.

C3 ~~6~~ ¹/~~16~~. (Thrice Amended) A method according to claim ~~10~~ ¹/~~10~~, wherein the composition is administered at a dosage so as to provide from about 0.1 to about 10 mg/kg of antibody or [antibody fragment] an antigen binding fragment of said antibody, based on the weight of the susceptible mammal.

C4 ~~7~~ ¹/~~17~~. (Amended) A method according to claim ~~10~~ ¹/~~10~~, wherein the composition is administered in an amount effective to [coat] block VLA-4 antigen on VLA-4 positive cells in the peripheral blood for a period of 1-14 days.

C5 ~~9~~ ¹/~~25~~. (Twice Amended) A method according to claim ~~10~~ ¹/~~10~~, wherein the composition comprises an antibody or an antigen binding fragment of [such] said antibody capable of binding to the α_4 subunit of VLA-4.

51